

## Patient Alert Card

### Apixaban JPI®

#### Apixaban

**Carry this card with you at all times.**

**Show this card to your pharmacist, dentist and any other healthcare professionals that treat you.**

#### Information for patients

- Take Apixaban JPI regularly as instructed. If you miss a dose, take it as soon as you remember and continue to follow your dosing schedule.
- Do not stop taking Apixaban JPI without talking to your doctor, as you are at risk of suffering from a stroke or other complications.
- Apixaban JPI helps to thin your blood. However, this may increase your risk of bleeding.
- Signs and symptoms of bleeding include bruising or bleeding under the skin, tar-coloured stools, blood in urine, nose-bleed, dizziness, tiredness, paleness or weakness, sudden severe headache, coughing up blood or vomiting blood.
- If the bleeding does not stop on its own, seek medical **attention immediately**.
- If you need surgery or any invasive procedure, inform your doctor that you are taking Apixaban JPI .

**I am under anticoagulation treatment with  
Apixaban JPI (apixaban) to prevent blood clots**

**Information for healthcare professionals**

- Apixaban JPI (apixaban) is an oral anticoagulant acting by direct selective inhibition of factor Xa.
- Apixaban JPI may increase the risk of bleeding. In case of major bleeding events, it should be stopped immediately.
- Treatment with Apixaban JPI does not require routine monitoring of exposure. A calibrated quantitative anti-Factor Xa assay may be useful in exceptional situations, e.g., overdose and emergency surgery (prothrombin time (PT), international normalised ratio(INR) and activated partial thromboplastin time (aPTT) clotting tests are not recommended) – see Summary of product characteristics (SPC).
- Agent to reverse the anti-factor Xa activity of apixaban is available.

**Please complete this section or ask your doctor to do it**

**Name:** \_\_\_\_\_

**Birth Date:** \_\_\_\_\_

**Indication:** \_\_\_\_\_

**Dose:** \_\_\_\_\_ **mg twice daily**

**Doctor's name:** \_\_\_\_\_

**Doctor's telephone:** \_\_\_\_\_

Any suspected adverse reactions to Apixaban JPI should be reported to Hikma Pharmaceuticals at the following email address: [SAPV@hikma.com](mailto:SAPV@hikma.com) and/or to Saudi Food and Drug Administration (SFDA) at the following contact details:

E-mail: [npc.drug@sfd.com](mailto:npc.drug@sfd.com),

Website: <https://ade.sfda.gov.sa>,

Call center number: 19999, or QR Code:

**Jazeera Pharmaceutical Industries**

Apixaban JPI 2.5 mg Film-coated Tablets, 20 Tablets

Apixaban JPI 2.5 mg Film-coated Tablets, 60 Tablets

Apixaban JPI 5 mg Film-coated Tablets, 60 Tablets

---

