

# **GUIDE FOR HEALTHCARE PROVIDERS CELLCEPT(MYCOPHENOLATE MOFETIL) RISK OF TERATOGENICITY**

## **INTRODUCTION**

This Guide has been designed to highlight the risks to babies associated with exposure to mycophenolate during pregnancy and to minimise the number of pregnancies during treatment with this teratogenic medicinal product.

Use this Guide during discussion with the patient and to address any questions or concerns the patient may have.

Although this Guide presents important information concerning the adverse pregnancy outcomes associated with mycophenolate, please consult the CellCept Summary of Product Characteristics (SmPC) for full information on mycophenolate.

## **PREGNANCY RISKS ASSOCIATED WITH MYCOPHENOLATE**

### **Preclinical evidence**

Mycophenolate is a powerful teratogen associated with an increased rate of spontaneous abortion and congenital malformation compared with other immunosuppressants. No specific mechanism of teratogenicity and mutagenicity has been identified. However, preclinical tests show fetal resorptions and malformations in rats and rabbits in the absence of maternal toxicity. Two genotoxicity assays indicated that mycophenolate has the potential to cause chromosomal damage at severely cytotoxic dose levels.

## **Clinical evidence in cases of maternal exposure**

A review of cumulative data found that around 45% to 49% of pregnancies in women exposed to mycophenolate resulted in spontaneous abortion, compared with reported frequencies of 12% to 33% in solid-organ transplantation patients treated with other immunosuppressants. The reported incidence of malformations in babies born to mothers exposed to mycophenolate during pregnancy is 23% to 27% compared with 4% to 5% in transplantation patients treated with other immunosuppressants, and 2% to 3% in the overall population.

Malformations associated with mycophenolate have included abnormalities of the ear, eye, and face; congenital heart disease including septal defects; polydactyly or syndactyly; tracheo-oesophageal malformations such as oesophageal atresia; effects on the nervous system such as spina bifida; and renal abnormalities.

## **Patients at risk of adverse pregnancy outcomes following exposure to mycophenolate include:**

- Pregnant patients.
- All female patients of childbearing potential (i.e. girls who have entered puberty and all women who have a uterus and have not passed through menopause).

## **Clinical evidence in cases of paternal exposure**

The limited clinical evidence available on paternally-exposed pregnancies does not indicate any increased risk of malformations or miscarriage following paternal exposure to mycophenolate.

**Mycophenolate is a powerful teratogen and may potentially be present in semen, however calculations on the amount that could potentially be transferred to a woman suggest it would be at a level unlikely to have an effect. Mycophenolate has been shown to be genotoxic in animal studies at concentrations exceeding the human therapeutic exposures only by small margins. Thus, the risk of genotoxic effects on sperm cells cannot be completely excluded.**

**As a precaution male patients and their female partners should be made aware of this potential risk and be recommended reliable contraceptive measures.**

## **PATIENT COUNSELLING**

**Before initiating or continuing treatment with mycophenolate, female and male patients must be educated about the increased risks of spontaneous abortion and congenital malformations associated with exposure to mycophenolate. You should ensure that women and men taking mycophenolate understand the risk of harm to the foetus, the need for effective contraception, and the need to immediately consult their physician if there is a possibility of pregnancy. The information you share in this discussion will be supported by the CellCept Guide for Patients and the Package Leaflet.**

## **In particular, you should:**

- **Counsel patients at risk to make sure they understand the risks and the measures required to minimise them.**
- **Provide female and male patients at risk with the CellCept Guide for Patients, and address any questions or concerns they might have.**
- **Explain the importance, methods, and timing of pregnancy tests prior and during treatment with mycophenolate.**
- **Provide counselling on the use of effective contraception prior to and during the entire duration of treatment with mycophenolate and for 6 weeks (female patients) or 90 days (male patients) after they stop taking mycophenolate.**
- **Advise patients taking mycophenolate that they must let you know in advance if they are considering becoming pregnant or fathering a child so that you can discuss possible treatment alternatives with them.**
- **Advise patients treated with mycophenolate not to donate blood during or for 6 weeks after stopping treatment. Male patients should not donate semen during therapy or for 90 days after stopping treatment.**
- **Advise patients that this medicine is for their own personal use, they should not give it to anyone else, and should return any unused medicine to their pharmacist at the end of treatment.**

## **PREGNANCY TESTING**

**Mycophenolate must not be used during pregnancy unless there is no suitable alternative to prevent transplant rejection.**

Before starting treatment with mycophenolate, women of childbearing potential should have two negative serum or urine pregnancy tests with sensitivity of at least 25 mIU/ml to exclude unintended exposure of the embryo to mycophenolate. It is recommended that the second test should be performed 8 to 10 days after the first one and immediately before starting mycophenolate mofetil. Pregnancy tests should be repeated as clinically required (e.g. after any gap in contraception is reported).

Results of all pregnancy tests should be discussed with the patient. Patients should be instructed to consult their physician immediately should pregnancy occur.

## **CONTRACEPTIVE REQUIREMENTS**

### **Women**

Mycophenolate is contraindicated in women of childbearing potential who are not using highly effective contraception. Because of the significant risks of spontaneous abortion and teratogenic potential of mycophenolate, women of childbearing potential must use at least one form of effective contraception before starting mycophenolate therapy, during therapy, and for 6 weeks after stopping the therapy; unless abstinence is the chosen method of contraception.

Two complementary forms of contraception are more effective and therefore preferred.

### **Men**

In the absence of sufficient data to exclude a risk of harm to the foetus, the following precautionary measures are recommended: sexually active male patients or their female partners are recommended to use reliable contraception during treatment of the male patient and for at least 90 days after cessation of mycophenolate mofetil.

## WHAT TO DO IF PREGNANCY OCCURS

You should base the course of action following exposure to mycophenolate during pregnancy on the individual patient's benefit- to – risk profile, and determine actions on a case by case basis through a discussion between the treating physician and the patient.

## CALL FOR REPORTING

- For full information on all possible adverse events please see the SPC
- Healthcare professionals are asked to report any suspected adverse reactions via the National reporting system that is provided below.
- Adverse reactions should also be reported to Roche Medical Information via the Company contact point that is provided below.
- In case of any adverse events – including any possible side effects not listed in the leaflet – or product complaints associated with the use of CellCept, please talk to the HCP or report the details in accordance with the national requirements via the national spontaneous reporting systems to:



The National Pharmacovigilance Centre  
Land Line: 19999.  
Fax: +966112057662  
Email: [npc.drug@sFDA.gov.sa](mailto:npc.drug@sFDA.gov.sa)



Roche Products Saudi Arabia L.L.C.  
Direct Tel. +966 12211 4618  
Mobile: +966 5678 44 692  
Email: [jeddah.drug\\_safety@roche.com](mailto:jeddah.drug_safety@roche.com)

Roche Products Saudi Arabia



This document has been reviewed and approved by The Saudi Food and Drug Authority (SFDA)